

Open Neighborhood Coloring of Line, Middle and Total Graphs of Some Graphs

R. Ezhilarasi

Ramanujan Institute for Advanced Study in Mathematics, University of Madras, Chennai - 600 005, India.

Abstract: In this paper, we determine the Open neighborhood chromatic number of line graph of Comb graph and double comb graph. Also, we obtain the open neighborhood chromatic number of the middle graph and total graph of path graph P_n , cycle graph C_n .

Keywords: simple graph, Open neighborhood coloring, line graph, middle graph, total graph. AMS Mathematics Subject Classification: 05C15, 05C62, 05C76

1 Introduction

All the graphs considered here are simple, connected and undirected graph $G = (V(G), E(G))$. For every vertex $a, b \in V(G)$, the edge connecting two vertices is denoted by $ab \in E(G)$. For all other standard concepts of graph theory, we see [1], [2], [5].

An open neighborhood k -coloring of a graph $G(V, E)$ is a k -coloring $f: V(G) \rightarrow \{1, 2, \dots, k\}$, $k \in \mathbb{Z}^+$ which admits the conditions such that for each $c \in V(G)$ and $\forall a, b \in N(c)$, $f(a) \neq f(b)$. The minimum value of k for which G admits an open neighborhood k -coloring is called the open neighborhood chromatic number of G and denoted by $\chi_{onc}(G)$.

[3], [4] Geetha K. N. introduced the notion and discussed the open neighborhood chromatic number of some graphs. Also, Adjacent vertex distinguishing total coloring of line and splitting graph of some graph and line graph of snake graph family has been obtained in the literature [6],

[7], [8].

The line graph $L(G)$ is the graph that represents the adjacencies between the edges of G . The vertex set of the middle graph $M(G)$ of the graph G is $V(G) \cup E(G)$ and in which two vertices are adjacent in $M(G)$ if and only if either they are adjacent edges of G or one is a vertex of G and the other is an edge incident with it. The total graph $T(G)$ of G has vertex set $V(G) \cup E(G)$, and edges joining all elements of this vertex set which are adjacent or incident in G .

Definition 1. The comb graph denoted by c^m is obtained from the path graph P_m with $\{v_1, v_2, \dots, v_m\}$ and the new vertices $\{u_1, u_2, \dots, u_m\}$ by joining the vertices $v_x u_x$ for $1 \leq x \leq m$.

$$V[c^m] = \left\{ \bigcup_{x=1}^m (u_x \cup v_x) \right\}$$

and

$$E[c^m] = \left\{ \left(\bigcup_{x=1}^{m-1} v_x v_{x+1} \right) \cup \left(\bigcup_{x=1}^m u_x v_x \right) \right\}$$

Here $d(v_1) = d(v_m) = 2$, $d(v_x) = 3$ for $2 \leq x \leq m-1$ and $d(u_x) = 1$ for $1 \leq x \leq m$.

Definition 2. The double comb graph denoted by Dc^m is obtained from the path graph P_m with $\{v_1, v_2, \dots, v_m\}$ and the new vertices $\{u_1, u_2, \dots, u_m\}$ and $\{w_1, w_2, \dots, w_m\}$ by joining the vertices $v_x u_x$ and $v_x w_x$ for

$1 \leq x \leq m$.

$$V[Dc^m] = \left\{ \bigcup_{x=1}^m u_x \cup v_x \cup w_x \right\}$$

and

$$E[Dc^m] = \left\{ \left(\bigcup_{x=1}^{m-1} v_x v_{x+1} \cup u_x v_x \cup v_x w_x \right) \right\}$$

Here $d(v_1) = d(v_m) = 3$, $d(v_x) = 4$ for $2 \leq x \leq m-1$ and $d(u_x) = d(w_x) = 1$ for $1 \leq x \leq m$.

2 Open neighborhood coloring of Line graph of comb and double comb graph

In this section, An open neighborhood coloring of line graph of comb graph c^m and double comb graph Dc^m are discussed.

Theorem 2.1. For Comb graph c^m , $\chi_{\text{onc}}(c^m) = 3$, for $m \geq 3$.

Proof. Let we denote $\{v_1, v_2, \dots, v_m, u_1, u_2, \dots, u_m\}$ as the vertices of c^m . Hence $|V(c^m)| = 2m$ and

$|E(c^m)| = 2m - 1$. Now we define $f: V(c^m) \rightarrow Z^+$ given by for $m \geq 3$,

$$\text{For } 1 \leq x \leq m, \quad f(v_x) = \begin{cases} \{1\}, & \text{if } x \equiv 1, 2 \pmod{4} \\ \{2\}, & \text{if } x \equiv 0, 3 \pmod{4} \end{cases}$$

$$\text{For } 1 \leq x \leq m, \quad f(u_x) = 3$$

It is easy to verify that f is an open neighborhood 3-coloring of c^m .

$$\therefore \chi_{\text{onc}}(c^m) = 3, \quad m \geq 3.$$

Hence the theorem. □

Theorem 2.2. For Double Comb graph Dc^m , $\chi_{\text{onc}}(Dc^m) = 4$, for $m \geq 3$.

Proof. Let we denote $\{v_1, v_2, \dots, v_m, u_1, u_2, \dots, u_m, w_1, w_2, \dots, w_m\}$ as the vertices of Dc^m . Hence

$|V(Dc^m)| = 3m$ and $|E(Dc^m)| = 3m - 1$. Now we define $f: V(Dc^m) \rightarrow Z^+$ given by for $m \geq 3$,

$$\text{For } 1 \leq x \leq m, \quad f(v_x) = \begin{cases} \{1\}, & \text{if } x \equiv 1, 2 \pmod{4} \\ \{2\}, & \text{if } x \equiv 0, 3 \pmod{4} \end{cases}$$

$$\text{For } 1 \leq x \leq m, \quad f(u_x) = 3 \text{ and } f(w_x) = 4.$$

It is easy to verify that f is an open neighborhood 4-coloring of Dc^m .

$$\therefore \chi_{\text{onc}}(Dc^m) = 4, \quad m \geq 3.$$

Hence the theorem. □

Theorem 2.3. For Line graph of Comb graph c^m , $\chi_{\text{onc}}(L(c^m)) = 5$, for $m \geq 4$.

Proof. From the definition (1), The Line graph of c^m is obtained by replacing all edges as vertices, we have $v_x v_{x+1} = s_x$, for $1 \leq x \leq m-1$, $u_x v_x = t_x$ for $1 \leq x \leq m$. Here s_x, t_x are the vertices of $L(c^m)$. Hence, the vertex set and edge set of $L(c^m)$ is given by

$$V[L(c^m)] = \left\{ \left(\bigcup_{x=1}^{m-1} s_x \right) \cup \left(\bigcup_{x=1}^m t_x \right) \right\}$$

$$E[L(c^m)] = \left\{ \left(\bigcup_{x=1}^{m-2} s_x s_{x+1} \right) \cup \left(\bigcup_{x=1}^{m-1} (s_x t_x \cup s_x t_{x+1}) \right) \right\}$$

Therefore $|V(c^m)| = 2m - 1$ and $|E(L(c^m))| = 3m - 4$. Now we define $f: V[L(c^m)] \rightarrow 5$ given by for $m \geq 4$,

$$\text{For } 1 \leq x \leq m-1, \quad f(s_x) = \begin{cases} \{1\}, & \text{if } x \equiv 1 \pmod{3} \\ \{2\}, & \text{if } x \equiv 2 \pmod{3} \\ \{3\}, & \text{if } x \equiv 0 \pmod{3} \end{cases}$$

$$\text{For } 1 \leq x \leq m, \quad f(t_x) = \begin{cases} \{4\}, & \text{if } x \equiv 1 \pmod{2} \\ \{5\}, & \text{if } x \equiv 0 \pmod{2} \end{cases}$$

It is easy to verify that f is an open neighborhood 5-coloring of $L(c^m)$.

$$\therefore \chi_{onc}(L(c^m)) = 5, \quad m \geq 4.$$

Hence the theorem. □

Theorem 2.4. For Line graph of double Comb graph Dc^m , $\chi_{onc}(L(Dc^m)) = 7$, for $m \geq 4$.

Proof. From the definition (2), The Line graph of Dc^m is obtained by replacing all edges as vertices, we have $v_x v_{x+1} = s_x$, for $1 \leq x \leq m-1$. $u_x v_x = r_x$ for $1 \leq x \leq m$. $v_x w_x = t_x$ for $1 \leq x \leq m$. Here s_x, r_x, t_x are the vertices of $L(Dc^m)$. Hence, the vertex set and edge set of $L(Dc^m)$ is given by

$$V[L(Dc^m)] = \left\{ \left(\bigcup_{x=1}^{m-1} s_x \right) \cup \left(\bigcup_{x=1}^m r_x \right) \cup \left(\bigcup_{x=1}^m t_x \right) \right\}$$

$$E[L(Dc^m)] = \left\{ \left(\bigcup_{x=1}^{m-2} (s_x s_{x+1}) \right) \cup \left(\bigcup_{x=1}^{m-1} (s_x r_x \cup s_x r_{x+1}) \right) \cup \left(\bigcup_{x=1}^{m-1} (s_x t_x \cup s_x t_{x+1}) \right) \cup \left(\bigcup_{x=1}^m (t_x r_x) \right) \right\}$$

Therefore $|V(L(Dc^m))| = 3m - 1$ and $|E(L(Dc^m))| = 6(m - 1)$. Now we define $f: V[L(Dc^m)] \rightarrow 7$ given by for $m \geq 4$,

$$\text{For } 1 \leq x \leq m-1, \quad f(s_x) = \begin{cases} \{1\}, & \text{if } x \equiv 1 \pmod{3} \\ \{2\}, & \text{if } x \equiv 2 \pmod{3} \\ \{3\}, & \text{if } x \equiv 0 \pmod{3} \end{cases}$$

$$\text{For } 1 \leq x \leq m, \quad f(r_x) = \begin{cases} \{6\}, & \text{if } x \equiv 1 \pmod{2} \\ \{4\}, & \text{if } x \equiv 0 \pmod{2} \end{cases}$$

$$\text{For } 1 \leq x \leq m, \quad f(t_x) = \begin{cases} \{7\}, & \text{if } x \equiv 1 \pmod{2} \\ \{5\}, & \text{if } x \equiv 0 \pmod{2} \end{cases}$$

It is easy to verify that f is an open neighborhood 7-coloring of $L(Dc^m)$.

$$\therefore \chi_{onc}(L(Dc^m)) = 7, \quad m \geq 4.$$

Hence the theorem. □

3 Open neighborhood coloring of Middle graph of Path and cycle graph

In this section, An open neighborhood coloring of middle graph of path P_n and cycle graph C_n are discussed.

Theorem 3.1. For middle graph of path graph $M(P_n)$, $\chi_{onc}(M(P_n)) = 5$, for $n \geq 4$

Proof. The middle graph of P_n is obtained from the path graph P_n with the vertices $\{v_1, v_2, \dots, v_n\}$ and let take the edges of P_n as vertices of $M(P_n)$, we have $v_x v_{x+1} = e_x$, for $1 \leq x \leq n-1$.

$$V[M(P_n)] = \left\{ \left(\bigcup_{x=1}^n v_x \right) \cup \left(\bigcup_{x=1}^{n-1} e_x \right) \right\}$$

and

$$E[M(P_n)] = \left\{ \left(\bigcup_{x=1}^{n-2} e_x e_{x+1} \right) \cup \left(\bigcup_{x=1}^{n-1} v_x e_x \right) \cup \left(\bigcup_{x=1}^{n-1} v_{x+1} e_x \right) \right\}$$

Now we define $f: V[M(P_n)] \rightarrow 5$ given by for $n \geq 4$,

$$\text{For } 1 \leq x \leq n-1, \quad f(e_x) = \begin{cases} \{1\}, & \text{if } x \equiv 1 \pmod{3} \\ \{2\}, & \text{if } x \equiv 2 \pmod{3} \\ \{3\}, & \text{if } x \equiv 0 \pmod{3} \end{cases}$$

$$\text{For } 1 \leq x \leq n, \quad f(v_x) = \begin{cases} \{4\}, & \text{if } x \equiv 1 \pmod{2} \\ \{5\}, & \text{if } x \equiv 0 \pmod{2} \end{cases}$$

It is easy to verify that f is an open neighborhood 5-coloring of $M(P_n)$.

$$\therefore \chi_{onc}(M(P_n)) = 5, \quad n \geq 4.$$

Hence the theorem. □

Theorem 3.2. For middle graph of cycle graph $M(C_n)$, for $n \geq 6$,

$$\chi_{onc}(M(C_n)) = \begin{cases} \{5\}, & \text{if } n \equiv 0, 5 \pmod{6} \\ \{6\}, & \text{otherwise} \end{cases}$$

Proof. The middle graph of C_n is obtained from C_n with the vertices $\{v_1, v_2, \dots, v_n\}$ and let take the edges of C_n as vertices of $M(C_n)$, we have $v_x v_{x+1} = e_x$, for $1 \leq x \leq n-1$ and $v_n v_1 = e_n$.

$$V[M(C_n)] = \left\{ \left(\bigcup_{x=1}^n v_x \cup e_x \right) \right\}$$

and

$$E[M(C_n)] = \left\{ \left(\bigcup_{x=1}^{n-1} e_x e_{x+1} \right) \cup \left(\bigcup_{x=1}^n v_x e_x \right) \cup \left(\bigcup_{x=1}^{n-1} v_{x+1} e_x \right) \cup (e_n e_1) \cup (v_1 e_n) \right\}$$

Then $|V(M(C_n))| = 2n$ and $|E(M(C_n))| = 3n$.

Define $f: V(M(C_n)) \rightarrow Z^+$ as follows.

Case-1. When $n \equiv 0 \pmod{6}$.

For $1 \leq x \leq n$,

$$f(e_x) = \begin{cases} 1, & \text{if } x \equiv 1 \pmod{3} \\ 2, & \text{if } x \equiv 2 \pmod{3} \\ 3, & \text{if } x \equiv 0 \pmod{3} \end{cases}$$

For $2 \leq x \leq n$,

$$f(v_x) = \begin{cases} 4, & \text{if } x \equiv 0 \pmod{2} \\ 5, & \text{if } x \equiv 1 \pmod{2} \end{cases} \text{ and } f(v_1) = 5.$$

$$\therefore \chi_{onc}(M(C_n)) = 5, \quad \text{for } n \equiv 0 \pmod{6}.$$

Case-2. When $n \equiv 1 \pmod{6}$.

For $1 \leq x \leq n-4$,

$$f(e_x) = \begin{cases} 1, & \text{if } x \equiv 1 \pmod{3} \\ 2, & \text{if } x \equiv 2 \pmod{3} \\ 3, & \text{if } x \equiv 0 \pmod{3} \end{cases}$$

$$f(e_{n-3}) = f(e_n) = 6, f(e_{n-2}) = 2, f(e_{n-1}) = 3$$

For $2 \leq x \leq n-2$,

$$f(v_x) = \begin{cases} 4, & \text{if } x \equiv 1 \pmod{2} \\ 5, & \text{if } x \equiv 0 \pmod{2} \end{cases}$$

$$\text{and } f(v_1) = 4, f(v_n) = 5, f(v_{n-1}) = 1.$$

$$\therefore \chi_{onc}(M(C_n)) = 6, \quad \text{for } n \equiv 1 \pmod{6}.$$

Case-3. When $n \equiv 2 \pmod{6}$.

For $1 \leq x \leq n-2$,

$$f(e_i) = \begin{cases} 1, & \text{if } x \equiv 1 \pmod{3} \\ 2, & \text{if } x \equiv 2 \pmod{3} \\ 3, & \text{if } x \equiv 0 \pmod{3} \end{cases}$$

$$f(e_{n-1}) = 4, \quad f(e_n) = 5$$

For $2 \leq x \leq n-3$,

$$f(v_x) = \begin{cases} 4, & \text{if } x \equiv 0 \pmod{2} \\ 5, & \text{if } x \equiv 1 \pmod{2} \end{cases}; f(v_x) = \begin{cases} 1, & \text{if } x = n-1 \\ 2, & \text{if } x = n \\ 3, & \text{if } x = 1 \end{cases}$$

$$\text{and } f(v_{n-2}) = 6.$$

$$\therefore \chi_{onc}(M(C_n)) = 6, \quad \text{for } n \equiv 2 \pmod{6}.$$

Case-4. When $n \equiv 3 \pmod{6}$.

For $1 \leq x \leq n$,

$$f(e_x) = \begin{cases} 1, & \text{if } x \equiv 1 \pmod{3} \\ 2, & \text{if } x \equiv 2 \pmod{3} \\ 3, & \text{if } x \equiv 0 \pmod{3} \end{cases}$$

For $2 \leq x \leq n$,

$$f(v_x) = \begin{cases} 4, & \text{if } x \equiv 0 \pmod{2} \\ 5, & \text{if } x \equiv 1 \pmod{2} \end{cases} \text{ and } f(v_1) = 6$$

$$\therefore \chi_{onc}(M(C_n)) = 6, \text{ for } n \equiv 3 \pmod{6}.$$

Case-5. When $n \equiv 4 \pmod{6}$.

For $1 \leq x \leq n-1$,

$$f(e_x) = \begin{cases} 1, & \text{if } x \equiv 1 \pmod{3} \\ 2, & \text{if } x \equiv 2 \pmod{3} \\ 3, & \text{if } x \equiv 0 \pmod{3} \end{cases} \text{ and } f(e_n) = 4$$

For $2 \leq x \leq n-2$,

$$f(v_x) = \begin{cases} 4, & \text{if } x \equiv 1 \pmod{2} \\ 5, & \text{if } x \equiv 0 \pmod{2} \end{cases} \text{ and } f(v_1) = f(v_{n-1}) = 6, f(v_n) = 5$$

$$\therefore \chi_{onc}(M(C_n)) = 6, \text{ for } n \equiv 4 \pmod{6}.$$

Case-6. When $n \equiv 5 \pmod{6}$.

For $1 \leq x \leq n-2$,

$$f(e_x) = \begin{cases} 1, & \text{if } x \equiv 1 \pmod{3} \\ 2, & \text{if } x \equiv 2 \pmod{3} \\ 3, & \text{if } x \equiv 0 \pmod{3} \end{cases}, \text{ and } f(e_{n-1}) = 4, f(e_n) = 5$$

For $2 \leq x \leq n-2$,

$$f(v_x) = \begin{cases} 4, & \text{if } x \equiv 0 \pmod{2} \\ 5, & \text{if } x \equiv 1 \pmod{2} \end{cases}; f(v_i) = \begin{cases} 1, & \text{if } x = n-1 \\ 2, & \text{if } x = n \\ 3, & \text{if } x = 1 \end{cases}$$

$$\therefore \chi_{onc}(M(C_n)) = 5, \text{ for } n \equiv 5 \pmod{6}.$$

It is easy to verify that f is an open neighborhood coloring of $M(P_n)$.

$$\therefore \chi_{onc}(M(C_n)) = \begin{cases} \{5\}, & \text{if } n \equiv 0, 5 \pmod{6} \\ \{6\}, & \text{otherwise} \end{cases}$$

Hence the theorem. □

4 Open neighborhood coloring of Total graph of Path and cycle graph

In this section, An open neighborhood coloring of total graph of path P_n and cycle graph C_n are discussed.

Theorem 4.1. For total graph of path graph $T(P_n)$, $\chi_{onc}(T(P_n)) = 5$, for $n \geq 4$

Proof. The total graph of P_n is obtained from the path graph P_n with the vertices $\{v_1, v_2, \dots, v_n\}$ and let take the edges of P_n as vertices of $T(P_n)$, we have $v_x v_{x+1} = e_x$, for $1 \leq x \leq n-1$.

$$V[T(P_n)] = \left\{ \left(\bigcup_{x=1}^n v_x \right) \cup \left(\bigcup_{x=1}^{n-1} e_x \right) \right\}$$

and

$$E[T(P_n)] = \left\{ \left(\bigcup_{x=1}^{n-1} v_x v_{x+1} \right) \cup \left(\bigcup_{x=1}^{n-2} e_x e_{x+1} \right) \cup \left(\bigcup_{x=1}^{n-1} v_x e_x \right) \cup \left(\bigcup_{x=1}^{n-1} v_{x+1} e_x \right) \right\}$$

Now we define $f: V[T(P_n)] \rightarrow 5$ given by for $n \geq 4$,

$$\text{For } 1 \leq x \leq n-1, \quad f(e_x) = \begin{cases} \{1\}, & \text{if } x \equiv 3 \pmod{5} \\ \{2\}, & \text{if } x \equiv 4 \pmod{5} \\ \{3\}, & \text{if } x \equiv 0 \pmod{5} \\ \{4\}, & \text{if } x \equiv 1 \pmod{5} \\ \{5\}, & \text{if } x \equiv 2 \pmod{5} \end{cases}$$

$$\text{For } 1 \leq x \leq n \text{ and } 1 \leq y \leq 4, \quad f(v_x) = y, \quad \text{if } x \equiv y \pmod{5}, f(v_{5x}) = 5$$

It is easy to verify that f is an open neighborhood 5-coloring of $T(P_n)$.

$$\therefore \chi_{onc}(T(P_n)) = 5, \quad n \geq 4.$$

Hence the theorem. □

Theorem 4.2. For total graph of cycle graph $T(C_n)$ for $n > 5$, $\chi_{onc}(T(C_n)) = \begin{cases} \{6\}, & \text{if } n \equiv 0, 2 \pmod{3} \\ \{7\}, & \text{otherwise} \end{cases}$

Proof. The total graph of C_n is obtained from C_n with the vertices $\{v_1, v_2, \dots, v_n\}$ and let take the edges of C_n as vertices of $T(C_n)$, we have $v_x v_{x+1} = e_x$, for $1 \leq x \leq n-1$ and $v_n v_1 = e_n$.

$$V[T(C_n)] = \left\{ \left(\bigcup_{x=1}^n v_i \cup e_x \right) \right\}$$

and

$$E[T(C_n)] = \left\{ \left(\bigcup_{x=1}^{n-1} e_x e_{x+1} \cup v_x v_{x+1} \right) \cup \left(\bigcup_{x=1}^n v_x e_x \right) \cup \left(\bigcup_{x=1}^{n-1} v_{x+1} e_x \right) \cup (e_n e_1) \cup (v_1 e_n) \cup (v_n v_1) \right\}$$

Then $|V(T(C_n))| = 2n$ and $|E(T(C_n))| = 4n$.

Define $f: V(T(C_n)) \rightarrow \mathbb{Z}^+$ as follows.

Case-1. When $n \equiv 0 \pmod{3}$.

$$\text{For } 1 \leq x \leq n, \quad f(e_x) = \begin{cases} 1, & \text{if } x \equiv 1 \pmod{3} \\ 2, & \text{if } x \equiv 2 \pmod{3} \\ 3, & \text{if } x \equiv 0 \pmod{3} \end{cases}$$

$$\text{For } 2 \leq x \leq n, \quad f(v_x) = \begin{cases} 4, & \text{if } x \equiv 2 \pmod{3} \\ 5, & \text{if } x \equiv 0 \pmod{3} \\ 6, & \text{if } x \equiv 1 \pmod{3} \end{cases} \quad \text{and } f(v_1) = 6.$$

$$\therefore \chi_{onc}(T(C_n)) = 6, \quad \text{for } n \equiv 0 \pmod{3}.$$

Case-2. When $n \equiv 1 \pmod{3}$.

$$\text{For } 1 \leq x \leq n-4,$$

$$f(e_x) = \begin{cases} 1, & \text{if } x \equiv 1 \pmod{3} \\ 2, & \text{if } x \equiv 2 \pmod{3} \\ 3, & \text{if } x \equiv 0 \pmod{3} \end{cases}$$

$$f(e_{n-3}) = 4, f(e_{n-2}) = 5, f(e_{n-1}) = 6, f(e_n) = f(v_{n-2}) = 7$$

$$\text{For } 2 \leq x \leq n-3,$$

$$f(v_x) = \begin{cases} 4, & \text{if } x \equiv 2 \pmod{3} \\ 5, & \text{if } x \equiv 0 \pmod{3} \\ 6, & \text{if } x \equiv 1 \pmod{3} \end{cases}$$

$$\text{and } f(v_1) = 3, f(v_n) = 2, f(v_{n-1}) = 1.$$

$$\therefore \chi_{onc}(T(C_n)) = 7, \text{ for } n \equiv 1 \pmod{3}.$$

Case-3. When $n \equiv 2 \pmod{3}$.

For $1 \leq x \leq n-2$,

$$f(e_x) = \begin{cases} 1, & \text{if } x \equiv 1 \pmod{3} \\ 2, & \text{if } x \equiv 2 \pmod{3} \\ 3, & \text{if } x \equiv 0 \pmod{3} \end{cases}$$

$$f(e_{n-1}) = 4, \quad f(e_n) = 5$$

For $2 \leq x \leq n-2$,

$$f(v_x) = \begin{cases} 4, & \text{if } x \equiv 2 \pmod{3} \\ 5, & \text{if } x \equiv 0 \pmod{3} \\ 6, & \text{if } x \equiv 1 \pmod{3} \end{cases}; f(v_x) = \begin{cases} 1, & \text{if } x = n-1 \\ 2, & \text{if } x = n \\ 3, & \text{if } x = 1 \end{cases}$$

$$\therefore \chi_{onc}(T(C_n)) = 6, \text{ for } n \equiv 2 \pmod{3}.$$

It is easy to verify that f is an open neighborhood coloring of $T(P_n)$.

$$\chi_{onc}(T(C_n)) = \begin{cases} \{6\}, & \text{if } n \equiv 0, 2 \pmod{3} \\ \{7\}, & \text{otherwise} \end{cases}$$

Hence the theorem. □

Conclusion.

In this paper, we have proved that the open neighborhood chromatic number of comb graph, double comb, line graph of comb, line graph of double comb graph. Also, we found the middle and total graph of path and cycle graph admits open neighborhood coloring conjecture.

References

- [1] N.L. Biggs, Algebraic Graph Theory, Cambridge University Press, 2nd edition, Cambridge, 1993.
- [2] Frank Harary, Graph Theory (Narosa publishing House. (2001)).
- [3] Geetha K N, Meera K N, Narahari N and Sooryanarayana B, Open Neighborhood Coloring of graphs, Int. J. Contemp. Math. Sciences, Vol.8 (14), (2013), pp.675-686.
- [4] Geetha K N, Meera K N, Narahari N and Sooryanarayana B, Open Neighborhood Coloring of prisms, J. Math. Fund. Sci., Vol.45 (3), (2013), pp. 245-262.
- [5] K.K.Parthasarathy, Basic Graph Theory, Tata McGraw Hill, 1994.
- [6] K.Thirusangu and R.Ezhilarasi "Adjacent-Vertex- Distinguishing total coloring of tensor product of graphs", American International Journal of Research in Science, Technology, Engineering & Mathematics, 24(1), September-November, 2018, pp. 52-61.
- [7] K.Thirusangu and R.Ezhilarasi "Adjacent Vertex Distinguishing total coloring of Line and splitting graph of some graph", Annals of Pure and Applied Mathematics, Vol 13, No.02, 2017, 173-183.
- [8] K.Thirusangu and R.Ezhilarasi "On Adjacent Vertex Distinguishing total coloring of Line graph of Snake graph family", International Journal of Pure and Applied Mathematics, Vol 113, No.06 - 2017, 261-269.