

Mangrove Tourism Suitability Index and Ecotourism Sustainability in the Waters of Talengen Bay, North Sulawesi Province, Indonesia

Elvi Siska Mandiangan^{1*}, Harsuko Rinwati², Abu Bakar Sambah³, Supriatna⁴

¹ Doctoral Program of Fisheries and Marine Sciences, Faculty of Fisheries and Marine Sciences, Jl. Veteran, Ketawanggede, District. Lowokwaru, Malang City, East Java 65145

² Faculty of Fisheries and Marine Sciences, Jl. Veteran, Ketawanggede, District. Lowokwaru, Malang City, East Java 65145

Abstract:- This study examines the potential of the mangrove ecosystem in Talengen Bay, North Sulawesi, Indonesia, for ecotourism development. Mangroves play a crucial role in environmental sustainability by supporting biodiversity, protecting coastlines, and providing habitat for aquatic species. The research aims to assess the tourism suitability index and the regional carrying capacity of the mangrove areas, crucial for sustainable tourism planning. The study identified six mangrove species, with *Rhizophora mucronata* being the most dominant. The Mangrove Tourism Suitability Index (MTSI) was calculated based on parameters such as mangrove thickness, density, species diversity, tidal influence, and associated biota. The results indicate that the Talengen Bay mangrove ecosystem is suitable for ecotourism, with an average MTSI score placing it in the "Suitable" category. Additionally, the carrying capacity for mangrove tourism activities was determined to be 105 visitors per day, ensuring minimal impact on the ecosystem. These findings provide valuable insights for policymakers and stakeholders to promote ecotourism while ensuring the conservation and sustainability of the mangrove ecosystem in Talengen Bay.

Keywords: Ecotourism, Tourism Suitability Index, Regional Carrying Capacity, Talengen Bay.

1. Introduction

The mangrove ecosystem has a very important role in supporting the sustainability of the surrounding environment (Abidin et al., 2021; Ruslan et al., 2022). Mangroves are coastal ecosystems that have a high level of productivity, so they are able to support the life of surrounding organisms and maintain ecosystem balance (K et al., 2024; Rahmadi et al., 2023). Apart from that, the mangrove ecosystem also acts as a habitat for various types of biotas, a spawning place for aquatic biota, a coastline protector, as well as a place for foraging and rearing of aquatic biota (Islamy & Hasan, 2020; Isoni et al., 2019, 2023).

The concept of ecotourism is defined as responsible travel to natural areas that preserves the environment, supports the well-being of local communities, and involves interpretation and education. (Chan et al., 2021; Huang et al., 2023). Ecotourism, which emphasizes mindful travel practices aimed at preserving the environment and supporting indigenous communities, has become a center of attention in assessing the suitability of Talengen Bay for tourism ventures (Satrya et al., 2023). In developing ecotourism, it is important to analyze the suitability of tourism and the carrying capacity of the area (Abdillah et al., 2020; Ariani et al., 2020). A study shows that the tourism suitability index, calculate the carrying capacity of the area, and identify ecotourism activities that can be carried out in mangrove areas (Ledheng et al., 2022; Vipriyanti et al., 2022). In addition, research also used analysis of the Tourism Suitability Index (TSI) and Ecotourism Area Carrying Capacity (ACC) for mangrove track exploration tourism (Nugraha et al., 2024). Indonesia has one of the largest mangrove forests in the world,

and North Sulawesi Province has extensive mangrove forests. Talengen Bay, Sangihe Islands District located in this province, stands out as a prime location for studying the mangrove ecosystem due to its relatively untouched condition and diverse wildlife. The potential to develop ecotourism in this region is seen as a dual opportunity: encouraging conservation efforts while providing economic benefits to local communities.

The development and use of mangrove ecosystems for ecotourism is an alternative sustainable development to overcome the problem of destructive use (Ferreira et al., 2022). However, in Talengen Bay, the implementation of mangrove ecotourism is not yet fully supported by the availability of physical data, especially information regarding the suitability and carrying capacity of the area. Therefore, there are several things that need attention and a broader and more comprehensive study is needed to determine the ability of the mangrove ecosystem to support all ecotourism activities being developed. This is because mangrove ecosystem resources are limited and vulnerable to both internal and external pressures which can reduce the quality and quantity of the ecosystem.

The results of this research are intended to provide valuable insights for policy makers, conservation activists and local stakeholders interested in promoting sustainable ecotourism in Talengen Bay. By understanding the current state and potential of mangrove ecosystems, effective strategies for conservation and sustainable tourism development can be formulated, thereby ensuring long-term benefits for the environment and local communities.

2. Material and Methods

Study Location

The research was conducted in Talengen Bay, North Sulawesi Province, Indonesia. This location is known for its extensive mangrove forests, providing an ideal setting for studying the suitability of mangrove tourism and its sustainability as an ecotourism destination (Figure 1).




Figure 1. Research location

Data collections

Data for this research was collected through a combination of field surveys, remote sensing, and secondary data sources. Primary data collection focused on mangrove types, mangrove density, mangrove thickness, associated biota objects and tracking length in the Talengen Bay Mangrove ecotourism area. Secondary data is tide data for the period October 2023 obtained from the Naval Hydro-Oceanography Center Institute (PUSHIDROSAL).

Field Surveys and Remote Sensing

Field surveys were conducted in Talengen Bay to assess the density and extent of mangrove forests using two main methods: transect sampling and GPS mapping. Transects, which are made perpendicular to the coastline, function as paths to create plots at regular intervals. In each plot, detailed data on the number of mangrove trees and species composition were recorded. At the same time, GPS devices were used to pinpoint the geographic coordinates of the plots, thereby facilitating the creation of accurate maps illustrating the distribution of mangrove forests throughout the bay.

Complementing these land-based efforts, remote sensing techniques are also being used to improve understanding of mangrove ecosystems. Landsat 8 satellite imagery and aerial photos underwent careful analysis, including image classification to distinguish mangrove areas from other land cover types. Additionally, temporal changes in mangrove forest cover were investigated through change detection methodology, which involves comparing images taken in different years. This multifaceted approach combines field survey data with remote sensing insights, providing a comprehensive understanding of the dynamics and spatial distribution of mangrove forests in Talengen Bay.

3. Data Analysis

Identify Types of Mangroves

The types of mangrove trees found at the research location were identified using an identification guidebook referring to (H Beyer E Kennedy, 2018; Harahab & Setiawan, 2017; Nelly et al., 2020).

Calculation of the density of mangrove species

To calculate the density of mangrove species, use the following formula (Murniasih et al., 2022):

$$D_i = \frac{n_i}{A}$$

Note:

D_i = Specific density (ind/m²)

n_i = Total number of i-type individuals

A = The total area of the sample observation area (m²)

Mangrove Thickness Calculation

Mangrove thickness analysis was carried out using the Google Earth application on the Asus Vivobook Core I9 13th Gen Laptop device. Mangrove thickness was measured at the research location where each transect location was located, namely measured from the outermost mangrove vegetation adjacent to the sea to the last (innermost) mangrove vegetation adjacent to land. The obtained mangrove thickness line length values are used to calculate the tourism suitability index value based on unpublished method (Tjong et al., 2021).

Tide Calculation

Tidal data obtained based on a specified time period, namely the highest and lowest tidal data, is further analyzed by adding up the two tidal values, which are then divided by two to obtain the average tidal value. The tidal data is used in the to get a score for the tidal parameters.

Biota Object Identification

Associated biota objects were found and documented at each research location where each transect was located using an Android camera, then these biota objects were grouped into the categories of fish, crabs, reptiles and birds. This data was further analyzed into a mangrove ecotourism suitability matrix to obtain a score for these parameters.

Mangrove Tourism Suitability Index (MTSI)

The mangrove tourism suitability index analysis is taken from the analysis developed published method (Hagger et al., 2022), where there are 5 parameters that must be calculated to get a score and category. The parameters considered include mangrove thickness, mangrove density, mangrove types, tides and associated biota objects. The formula used is as follows:

$$IKW = \sum_{i=1}^n Bi \times Xi$$

Note:

n = Number of Conformity Parameters

Bi = Parameter Weight i-th

Si = Score of the i-th Parameter

After the TSI value is obtained from each transect, the value is categorized according to the value provisions set by Yulianda (2019):

Very suitable = > 2,5

Suitable = 2,0 < TSI < 2,5

Not Appropriate = 1 < TSI < 2,0

Very Inappropriate = < 1

Regional Carrying Capacity

The area carrying capacity value (ACC) is needed to determine the maximum number of visitors that can be accommodated by the tourist area without causing disturbance to nature and humans. The ACC calculation uses the formula (Yulianda, 2019) as follows:

$$ACC = K \times \left(\frac{Lp}{Lt} \right) \times \left(\frac{Wt}{Wp} \right)$$

Note:

- ACC = Area Carrying Capacity (people/day)
- K = Ecological potential of visitors per unit area (persons)
- Lp = The area (m²) or length of the area (m) used
- Lt = Area units for specific categories (m² or m)
- Wt = Time provided by the area for tourist activities in 1 day (hours)
- Wp = Time spent by visitors for each activity (hours)

4. Results and Discussion

Results of Mangrove Type Identification

Based on the research results, 6 species were found, namely: *Rhizophora mucronata*, *Rhizophora apiculata*, *Bruguiera gymnorhiza*, *Sonneratia alba*, *Avicennia marina* and *Nypa fruticans* and dominated by *Rhizophora mucronata*. Mantehage Island, Bunaken National Park, Manado City, namely 20 species of mangroves (*Acanthus ilicifolius*, *Avicennia marina*, *Camptostemon philippinense*, *Nypa fruticans*, *Lumnitzera racemosa* Willd, *Cockroach Excoecaria*, *Sonneratia alba*, *Sonneratia ovata*, *Heritiera littoralis* Dryand , *Xylocarpus granatum*, *Aegiceras corniculatum*, *Acrostichum speciosum* Willd, *Bruguiera cylindrica*, *Bruguiera gymnorhiza*, *Bruguiera parviflora*, *Ceriops tagal*, *Rhizophora apiculata*, *Rhizophora mucronata*, *Rhizophora stylosa*, and *Scyphiphora hydrophyllacea*). According to Rumengan, et al. (2018), stated that differences in mangrove species in an area are influenced by physical and chemical factors in the water.

The diversity of mangrove types in the area is considered one of the key attractions for visitors, who engage in tours and educational activities related to the mangrove ecosystem (Waters & Gibbons, 2004). This diversity is viewed as an added value for the development of ecotourism in the Talengen Bay area. According to published method, the diversity of mangrove types in an area is recognized as an important value in supporting the management activities of a tourist area and increasing visitor attraction (Ewaldo et al., 2023; Getzner & Islam, 2020). A large number of mangrove species is believed to support overall biota diversity, thereby enhancing the ecological richness and appeal of the area (Irwansyah et al., 2022; Sukuryadi et al., 2020). This diversity is perceived to contribute significantly to the educational and recreational experiences of visitors, promoting a deeper understanding and appreciation of mangrove ecosystems (Sariaslan & Langer, 2021). Consequently, the management and conservation of these diverse mangrove species are seen as crucial for the sustainable development of ecotourism in Talengen Bay.

Calculation Results of Mangrove Type Density

Based on the research results, the density of mangrove species can be seen in Table 1.

Table 1. Calculation of Mangrove Species Density

Species	T1	T2	T3	T4	T5
<i>Rhizophora mucronata</i>	0.07	0.09	0.10	0.09	0.07
<i>Rhizophora apiculata</i>	0.03	0.03	0.03	0.02	0.00
<i>Bruguiera gymnorhiza</i>	0.01	0.01	0.00	0.01	0.01
<i>Sonneratia alba</i>	0.01	0.00	0.01	0.00	0.00
<i>Avicennia marina</i>	0.01	0.00	0.00	0.00	0.00
TOTAL	0.124	0.114	0.13	0.12	0.09

Based on the type of mangrove, *R. apiculata* is shown to have the highest species density in each transect, with a range value of 0.07–0.10 individuals per 500 m² and an average value of 0.08. The smallest density values for mangrove species are reported for *Avicennia marina* and *Bruguiera gymnorhiza*, with respective range values of 0–0.01 individuals per 500 m² and 0–0.4 individuals per 100 m², and average values of 0.02 and 0.01 individuals per 100 m². Studies have indicated that *Rhizophora* mangrove species tend to be dominant and abundant in mangrove communities, particularly due to their good adaptability to the typical environmental conditions in mangrove forests (Fazlioglu & Chen, 2020; Khan et al., 2024). The relatively high density in *Rhizophora* mangroves is attributed to a combination of the species' adaptability, complex root structure, interaction with the nekton community (Friess et al., 2021), as well as environmental factors that support the growth and proliferation of this

species in the mangrove ecosystem. This dominance and adaptability are considered crucial for maintaining the structure and function of mangrove ecosystems, thus enhancing their ecological value and resilience.

Mangrove Thickness Results

The thickness of the mangroves in the various transects is recorded as follows: 238 m in transect 1, 163 m in transect 2, 204 m in transect 3, 287 m in transect 4, and 132 m in transect 5. From this data, the average thickness of mangroves at the research location is calculated to be 204.8 m. This average thickness reflects the overall density and health of the mangrove ecosystem in the area. Mangrove conditions can serve as a significant attraction for tourists due to their aesthetic appeal (Murniasih et al., 2022).

The denser the mangrove forest, the more appealing it is considered to be for tourists and visitors. Additionally, dense mangrove forests are known to provide better habitat and protection for various species, contributing to biodiversity (Borsje et al., 2011). The thick mangrove stands also play a critical role in coastal protection, reducing the impact of waves and storm surges, which is another aspect that enhances their attractiveness for eco-tourists interested in the ecological and protective functions of mangroves (Borsje et al., 2011; De Silva & Amarasinghe, 2023; Friess et al., 2021). This combination of aesthetic and ecological benefits makes dense mangrove forests highly valuable for sustainable tourism development.

Tide Calculation Results

In October 2023, an analysis of tidal data from the Indonesian Navy's Hydro Oceanography Center (PUSHIDROSAL) revealed that the average tidal conditions around Talengen Bay stood at 1.89 meters. This figure represents an optimal value for mangrove tourism activities, as it does not surpass the height of the trail at this particular tourist destination. Consequently, tourists can comfortably engage in tracking and exploration activities.

The height of sea tides serves as a crucial indicator of visitor comfort during such endeavors, as noted by published article (Shahrin et al., 2024; Su et al., 2021). This finding underscores the importance of considering tidal patterns in designing and managing tourist experiences within coastal ecosystems. Moreover, it emphasizes the need for ongoing monitoring and research to ensure sustainable tourism practices that align with natural environmental dynamics.

Biota Object Identification Results

The research findings on the biotic components of the mangrove ecosystem in Talengen Bay reveal the presence of six distinct groups of biota, including fish, shrimp, crabs, mollusks, reptiles, and birds. The diversity of these biotic groups within the mangrove association is intricately linked to the thickness of the mangrove itself (Jacquot et al., 2023).

It is elucidated that as the thickness of the mangrove increases, a corresponding rise in the diversity of associated biota is observed. Conversely, a decrease in mangrove thickness is associated with a reduction in the variety of associated biota (Palit et al., 2022). This assertion underscores the significant role of mangrove structure and density in shaping the ecological dynamics of these coastal habitats, as highlighted by Sari's findings. Furthermore, it emphasizes the importance of preserving and managing mangrove ecosystems to sustain biodiversity and support ecotourism activities in regions like Talengen Bay.

Mangrove Tourism Suitability Index (MTSI)

The suitability of mangrove tourism considers five parameters, including mangrove thickness, mangrove density, mangrove type, tides and biota objects (Ewaldo et al., 2023; Ledheng et al., 2022). The results of the calculation of the mangrove tourism suitability index for each transect, namely transect 1 is 2.63, transect 2 is 2.25, transect 3 is 2.63, transect 4 is 2.63 and transect 5 is 1.85 while the average value The mangrove tourism suitability index for the five transects is 2.398 and this value means that the Talengen Mangrove Ecotourism area is categorized as suitable based on the published provisions.

The ACC assessment is said to be suitable if it has a value of <2.0. The ACC assessment shows that the Talengen Bay mangrove ecosystem is suitable for use as mangrove ecotourism because the calculated parameters have met the requirements and are suitable for development as an ecotourism attraction. such as mangrove types, mangrove thickness, mangrove density, tides and biota objects.

Table 2. Mangrove Tourism Suitability Index (MTSI) result

Parameters	Value weight	Transect 1			Transect 2			Transect 3			Transect 4			Transect 5		
		C	S	C*S	C	S	C*S	C	S	C*S	C	S	C*S	C	S	C*S
Mangrove thickness	0.38	>200-500 (m)	3	1.14	50-200 (m)	2	0.76	>200-500 (m)	3	1.14	>200-500 (m)	3	1.14	50-200 (m)	2	0.76
Mangrove type density	0.25	>10-15 (100 m ²)	2	0.5	>10-15 (100 m ²)	2	0.5	>10-15 (100 m ²)	2	0.5	>10-15 (100 m ²)	2	0.5	5-10 (100 m ²)	1	0.25
Type of Mangrove	0.15	>5 (types)	3	0.45	>5 (types)	3	0.45	>5 (types)	3	0.45	>5 (types)	3	0.45	>5 (types)	2	0.3
Tidal	0.12	1.89	2	0.24	1.89	2	0.24	1.89	2	0.24	1.89	2	0.24	1.89	2	0.24
Biota objects	0.1	Fish, crabs, mollusks, birds	3	0.3	Fish, crabs, mollusks, birds	3	0.3	Fish, crabs, mollusks, birds	3	0.3	Fish, crabs, mollusks, birds	3	0.3	Fish, crabs, mollusks, birds	3	0.3
ACC				2.63			2.25			2.63			2.63			1.85
Category		Sufficiently Sustainable			Sufficiently Sustainable			Sufficiently Sustainable			Sufficiently Sustainable			Less Sustainable		
Average ACC		2.398 (Sufficiently Sustainable)														
Category (ACC) Yulianda 2019		Appropriate = 2.0 < ACC < 2.5 Sufficiently Sustainable														

Regional Carrying Capacity (RCC)

The calculation of area carrying capacity in the Talengen Bay area involves assessing the length of the managed tracking routes by the local community. Utilizing Landsat Image 8 calculations, it was determined that the total area of mangrove forest in Talengen Bay spans 440,000 square meters, with the tracking route covering 4,500 square meters (Lp). Additionally, measurements of the tracking length revealed a distance of 300 meters (Lt). The time allocated for touring the attraction within a day is set at 10 hours (07:00-17:00 WITA) (Wt), with each visitor spending approximately 2 hours navigating the mangrove (Wp). Consequently, the area carrying capacity for Talengen Bay's mangrove ecotourism is estimated at 105 people per day. This calculation accounts for existing facilities such as gazebos, toilets, culinary offerings, washing areas, waste disposal facilities, and picturesque photography spots, with an entrance ticket selling at Rp. 5,000 per person.

Table 3. Area Carrying Capacity Parameters

Parameter	Obtained Data	Area Carrying Capacity
Ecological potential of visitors per unit area (persons)	1	105 persons/day
Usable area (m ²)	4500 m ²	
Area unit (area or length) for specific categories (m ² or m)	150 m	
Time provided for activities in one day (hours)	7	
Time spent by visitors for each activity (hours)	2	

The daily visitor count derived from area carrying capacity calculations serves as a crucial benchmark for regulating tourist influx, thereby mitigating adverse impacts on the ecological sustainability of the ecotourism site (Jamin & Rahmafritria, 2022). This practice aligns with sustainable tourism principles by ensuring that visitor numbers are maintained at levels that the ecosystem can support without degradation. By implementing such

measures, stakeholders can safeguard the long-term viability of Talengen Bay's mangrove ecotourism while simultaneously offering memorable experiences to visitors.

5. Conclusion

The types of mangroves found in the Talengen Bay mangrove ecosystem area include *Rhizophora mucronata*, *Rhizophora apiculata*, *Bruguiera gymnorhiza*, *Sonneratia alba*, *Avicennia marina* and *Nypa fruticans* and are dominated by *Rhizophora mucronata*. Based on the Tourism Suitability Index (TSI) for mangrove ecotourism in the Talengen Bay mangrove ecotourism area, it is included in the Suitable category with an average ACC value of 2.398. The carrying capacity of the mangrove forest area for tracking is 105 people/day,

Acknowledgement

Authors thanks to Provincial Government of North Sulawesi for funding and scholarships.

References

- [1] Abdillah, S., Harahab, N., Primyastanto, M., & Semedi, B. (2020). Analysis of suitability and carrying capacity of mangrove ecosystem for ecotourism in Lembar Village, West Lombok District, Indonesia. *Biodiversitas*, 21(2). <https://doi.org/10.13057/biodiv/d210222>
- [2] Abidin, Z., Setiawan, B., Muhaimin, A. W., & Shinta, A. (2021). The role of coastal biodiversity conservation on sustainability and environmental awareness in mangrove ecosystem of southern Malang, Indonesia. *Biodiversitas*, 22(2). <https://doi.org/10.13057/biodiv/d220217>
- [3] Ariani, F., Rahimah, I., Sipahutar, R., Yanti, E. S., & Paradini, A. (2020). Suitability and carrying capacity analyses for marine ecotourism development of the Medang Deras Subdistrict, Batubara district, province of North Sumatera. *IOP Conference Series. Earth and Environmental Science*, 584(1), 12010. <https://doi.org/10.1088/1755-1315/584/1/012010>
- [4] Borsje, B. W., Van Wesenbeeck, B. K., Dekker, F., Paalvast, P., Bouma, T. J., Van Katwijk, M. M., & De Vries, M. B. (2011). How ecological engineering can serve in coastal protection. *Ecological Engineering*, 37(2), 113–122. <https://doi.org/10.1016/j.ecoleng.2010.11.027>
- [5] Chan, J. K. L., Marzuki, K. M., & Mohtar, T. M. (2021). Local community participation and responsible tourism practices in ecotourism Destination: a case of Lower Kinabatangan, Sabah. *Sustainability*, 13(23), 13302. <https://doi.org/10.3390/su132313302>
- [6] De Silva, W., & Amarasinghe, M. D. (2023). Coastal protection function of mangrove ecosystems: a case study from Sri Lanka. *Journal of Coastal Conservation*, 27(6). <https://doi.org/10.1007/s11852-023-00990-8>
- [7] Ewaldo, N. K., Kurniasa, N. M., & Takarina, N. N. D. (2023). Suitability of mangrove forests for sustainable tourism. *Advances in Tropical Biodiversity and Environmental Sciences*, 7(3), 93–98. <https://doi.org/10.24843/atbes.2023.v07.i03.p02>
- [8] Ferreira, A. C., Borges, R., & De Lacerda, L. D. (2022). Can sustainable development save mangroves? *Sustainability*, 14(3), 1263. <https://doi.org/10.3390/su14031263>
- [9] Friess, D. A., Chua, S. C., Jaafar, Z., Krauss, K. W., & Yando, E. S. (2021). Mangroves and people: Impacts and interactions. *Estuarine, Coastal and Shelf Science*, 248, 107155. <https://doi.org/10.1016/j.ecss.2020.107155>
- [10] Getzner, M., & Islam, M. S. (2020). Ecosystem Services of Mangrove Forests: Results of a Meta-Analysis of Economic Values. *International Journal of Environmental Research and Public Health*. <https://doi.org/10.3390/ijerph17165830>
- [11] H Beyer E Kennedy, M. B. C. C. J. C. E. D. O. H.-G. (2018). Risk-sensitive planning for conserving coral reefs under rapid climate change. *Conservation Letters*, 6, 11.
- [12] Hagger, V., Worthington, T. A., Lovelock, C. E., Adame, M. F., Amano, T., Brown, B. M., Friess, D. A., Landis, E., Mumby, P. J., Morrison, T. H., O'Brien, K. R., Wilson, K. A., Zganjar, C., & Saunders, M. I. (2022). Drivers of global mangrove loss and gain in social-ecological systems. *Nature Communications*, 13(1). <https://doi.org/10.1038/s41467-022-33962-x>

- [13] Harahab, N., & Setiawan, S. (2017). Suitability index of mangrove ecotourism in malang regency. *Ecofom (Economic and Social of Fisheries and Marine)/ECSOFiM (Economic and Social of Fisheries and Marine)*, 004(02), 153–165. <https://doi.org/10.21776/ub.ecsofim.2017.004.02.05>
- [14] Huang, C.-C., Li, S.-P., Chan, Y.-K., Hsieh, M.-Y., & Lai, J.-C. M. (2023). Empirical Research on the Sustainable Development of Ecotourism with Environmental Education Concepts. *Sustainability*, 15(13), 10307. <https://doi.org/10.3390/su151310307>
- [15] Irwansyah, R. M., Azzahra, S. I. N., Darmastuti, S. A., Ramadhandi, A. R., Firdaus, O., Daeni, F., Safitri, N., Fajri, O. P. A., Nugroho, G. D., Naim, D. M., & Setyawan, A. D. (2022). Crab diversity and crab potential as support ecotourism in Teleng Ria, Grindulu and Siwil Beach, Pacitan, East Java, Indonesia. *International Journal of Bonorowo Wetlands*, 11(2). <https://doi.org/10.13057/bonorowo/w110204>
- [16] Islamy, R. A., & Hasan, V. (2020). Checklist of mangrove snails (Mollusca: Gastropoda) in South Coast of Pamekasan, Madura Island, East Java, Indonesia. *Biodiversitas*, 21(7). <https://doi.org/10.13057/biodiv/d210733>
- [17] Isoni, W., Islamy, R. A., Musa, M. Z., & Wijanarko, P. (2019). Short Communication: Species composition and density of mangrove forest in Kedawang Village, Pasuruan, East Java, Indonesia. *Biodiversitas*. <https://doi.org/10.13057/biodiv/d200626>
- [18] Isoni, W., Sari, P. D. W., Sari, L. A., Daniel, K., South, J., Islamy, R. A., Wirabuana, P. Y. A. P., & Hasan, V. (2023). Checklist of mangrove snails (Gastropoda: Mollusca) on the coast of Lamongan District, East Java, Indonesia. *Biodiversitas*, 24(3). <https://doi.org/10.13057/biodiv/d240341>
- [19] Jacquot, M. P., Nordström, M. C., De Wever, L., Ka, R. N., Ka, S., Garrec, V. Le, Raffray, J., Sadio, O., Diouf, M., Grall, J., De Morais, L. T., & Loc'h, F. Le. (2023). Human activities and environmental variables drive infaunal community structure and functioning in West African mangroves. *Estuarine, Coastal and Shelf Science*, 293, 108481. <https://doi.org/10.1016/j.ecss.2023.108481>
- [20] Jamin, A., & Rahmafritria, F. (2022). Visitor management concept through carrying capacity analysis in forest recreation. *Journal of Indonesian Tourism, Hospitality and Recreation*, 5(1), 1–12. <https://doi.org/10.17509/jithor.v5i1.46821>
- [21] K, A., K, H. P., K, S. V, P, B., Muhammed, J., & Augustine, A. (2024). Mangroves in environmental engineering: Harnessing the multifunctional potential of Nature's coastal architects for sustainable ecosystem management. *Results in Engineering*, 21, 101765. <https://doi.org/10.1016/j.rineng.2024.101765>
- [22] Ledheng, L., Dewi, N. P. Y. A., & Hale, M. (2022). The Analysis Of Combatibility And Support Capacity In The Mangrove Tourism Area At Susuk Pool, Dualaus Village, Belu District, East Nusa Tenggara Province. *Media Konservasi*, 27(3), 91–100. <https://doi.org/10.29244/medkon.27.3.91-100>
- [23] Murniasih, S., Hendarto, E., & Hilmi, E. (2022). The Mangrove Density, Diversity, and Environmental Factors as Important Variables to Support the Conservation Program of Essential Ecosystem Area in Muara Kali Ijo, Pantai Ayah, Kebumen. In *Jurnal Sylva Lestari*. <https://doi.org/10.23960/jsl.v10i3.596>
- [24] Nelly, C., Rasnovi, S., & Zumaidar, Z. (2020). Mangrove ecosystem suitability for Ecotourism Management Recommendation in Iboih Village – Sabang. *E3S Web of Conferences*, 151, 1060. <https://doi.org/10.1051/e3sconf/202015101060>
- [25] Nugraha, W. A., Ambariyanto, A., & Insafitri, I. (2024). The mangrove ecosystems on the north coast of Bangkalan, Indonesia, and its potential for mangrove ecotourism. *Biodiversitas*, 25(2). <https://doi.org/10.13057/biodiv/d250246>
- [26] Palit, K., Rath, S., Chatterjee, S., & Das, S. (2022). Microbial diversity and ecological interactions of microorganisms in the mangrove ecosystem: Threats, vulnerability, and adaptations. *Environmental Science and Pollution Research International*, 29(22), 32467–32512. <https://doi.org/10.1007/s11356-022-19048-7>
- [27] Rahmadi, M. T., Yuniastuti, E., Suciani, A., Harefa, M. S., Persada, A. Y., & Tuhono, E. (2023). Threats to mangrove ecosystems and their impact on coastal biodiversity: A study on mangrove management in Langsa City. *Indonesian Journal of Earth Sciences/Indonesian Journal Of Earth Sciences*, 3(2), A627. <https://doi.org/10.52562/injoes.2023.627>

-
- [28] Ruslan, N. F. N., Goh, H. C., Hattam, C., Edwards-Jones, A., & Moh, H. H. (2022). Mangrove ecosystem services: Contribution to the well-being of the coastal communities in Klang Islands. *Marine Policy*, *144*, 105222. <https://doi.org/10.1016/j.marpol.2022.105222>
- [29] Sariaslan, N., & Langer, M. R. (2021). Atypical, high-diversity assemblages of foraminifera in a mangrove estuary in northern Brazil. *Biogeosciences*, *18*(13), 4073–4090. <https://doi.org/10.5194/bg-18-4073-2021>
- [30] Satrya, I. D. G., Kaihatu, T. S., Budidharmanto, L. P., Karya, D. F., & Rusadi, N. W. P. (2023). The role of ecotourism in preserving environmental awareness, cultural and natural attractiveness for promoting local communities in Bali, Indonesia. *Journal of Eastern European and Central Asian Research*, *10*(7), 1063–1075. <https://doi.org/10.15549/jeeecar.v10i7.1386>
- [31] Shahrin, F., Hassan, Z. A., & Zahit, R. A. (2024). Re-visit project success: evaluation of visitor perspectives in the Sungai Melaka flood mitigation project. *Journal of Engineering, Design and Technology*. <https://doi.org/10.1108/jedt-07-2023-0285>
- [32] Su, J., Friess, D. A., & Gasparatos, A. (2021). A Meta-Analysis of the Ecological and Economic Outcomes of Mangrove Restoration. *Nature Communications*. <https://doi.org/10.1038/s41467-021-25349-1>
- [33] Sukuryadi, N., Harahab, N., Primyastanto, M., & Semedi, B. (2020). Collaborative-based mangrove ecosystem management model for the development of marine ecotourism in Lembar Bay, Lombok, Indonesia. *Environment, Development and Sustainability*, *23*(5), 6838–6868. <https://doi.org/10.1007/s10668-020-00895-8>
- [34] Tjiong, G. P., Supriatna, J., Takarina, N. D., & Tambunan, R. P. (2021). Mangrove diversity and suitability assessments for ecotourism in Cimalaya Wetan Coast, Karawang District, Indonesia. *Biodiversitas*, *22*(2). <https://doi.org/10.13057/biodiv/d220234>
- [35] Vipriyanti, N. U., Semadi, I. G. N. M. D., & Fauzi, A. (2022). Developing mangrove ecotourism in Nusa Penida Sacred Island, Bali, Indonesia. *Environment, Development and Sustainability*, *26*(1), 535–548. <https://doi.org/10.1007/s10668-022-02721-9>
- [36] Waters, S. H., & Gibbons, A. S. (2004). Design languages, notation systems, and instructional technology: A case study. *Educational Technology Research and Development*, *52*(2), 57–68. <https://doi.org/10.1007/BF02504839>